



Loomis Basin Equine Medical Center, Inc.

3901 Sierra College Blvd. - P.O. Box 2059 Loomis, CA 95650
(916) 652-7645 Fax (916) 652-7662 www.loomisbasinlargeanimal.com

Recommended Adult* Equine Preventative Vaccination Program (1/08) (*Horses 1 year of Age or Older)

VACCINE	ADMINISTERED <i>IM - Intramuscular</i> <i>IN - Intranasal</i>	FREQUENCY
Tetanus/Sleeping Sickness ¹	IM	Annually ⁴
Tetanus/Sleeping Sickness/West Nile Virus Combination	IM	Annually ⁴
Rabies	IM	Annually ⁴
Flu/Rhino - Killed Virus	IM or IN	Every 6 months
Rhino EHV 1 Modified Live Virus	IM	Every 3 months ³
Influenza ⁶	IM	Every 6 months
Strangles (IN) ²	IN	Annually ⁴
West Nile	IM	Twice Annually ^{4,5}

¹ In case of a penetrating wound, Tetanus Toxoid should be given if greater than 6 months since vaccination. If no prior Tetanus Toxoid can be established, consider Tetanus Antitoxin.

² We recommend the use of this vaccine only where there is known or anticipated exposure.

³ Where exposure may be high, we recommend this vaccine be given every 3 months. Start this series with an initial vaccine and a booster at 3 weeks. This vaccine may have some efficacy against the neurological form of Rhinopneumonitis. See Note 6 below regarding influenza vaccine.

⁴ Previously unvaccinated adults should be boosted 3-4 weeks after their initial vaccine. If using Tetanus/Sleeping Sickness/West Nile Combo, booster West Nile alone 6 months later.

⁵ If your horse was fully vaccinated last year or if the primary vaccination series was completed this year, follow up with a booster every six months. Try to time the summer booster for late June or early July due to high incidence of the disease in late summer.

⁶ If using the Rhino EHV-1 Modified Live Virus, an influenza vaccine must be given separately. Can be given separately or in combination with Killed Rhino (EHV-1) one month before foaling.